



INVESTOR IN PEOPLE

15
The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ


CERTIFIED COPY OF PRIORITY DOCUMENT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *AmBrewer*

Dated 31 July 2003

THIS PAGE BLANK (USPTO)

23 SEP 1999

LONDON

24SEP99 E479103-3 002000
F01/7700 0.00 - 9922561.7

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road

Newport

Gwent NP9 1RH

GWS\DJC\21091-01

1. Your reference

2. Patent application number

(The Patent Office will fill in this part)

23 SEP 1999

9922561.7

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Microbiological Research Authority

CAMR

Porton Down

Salisbury

Wiltshire SP4 0JG

United Kingdom

6566160001

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

4. Title of the invention

Neisserial Vaccine Compositions and Methods

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

MATHYS & SQUIRE

100 Grays Inn Road

London WC1X 8AL

1081001 ✓

Patents ADP number (if you know it)

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

a) any applicant named in part 3 is not an inventor, or

b) there is an inventor who is not named as an applicant, or

c) any named applicant is a corporate body.

See note (d))

YES

CERTIFICATE
ISSUED

2.3.00.

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 21

Claim(s) 6

Abstract -

Drawing(s)

8 + 6

10. If you are also filing any of the following, state how many against each item.

Priority documents -

Translations of priority documents -

Statement of inventorship and right to grant of a patent (Patents Form 7/77) -

Request for preliminary examination and search (Patents Form 9/77) -

Request for substantive examination (Patents Form 10/77) -

Any other documents (please specify) -

11.

I/We request the grant of a patent on the basis of this application.

Signature

Mathy, & Squire

Date 23 September 1999

12. Name and daytime telephone number of person to contact in the United Kingdom

George W Schlich

0171 830 0000

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

NEISSERIAL VACCINE COMPOSITIONS AND METHODS

The present invention relates to vaccines and methods for preparing vaccines that stimulate an immune response. In particular, the present invention relates to vaccines that provide broad spectrum protective immunity to microbial infection.

Infection by pathogenic organisms is one of the major causes of chronic and acute disease. In particular, infection resulting from microbial sources - such as bacteria, viruses and protozoans - continue to claim millions of lives worldwide. With microbial species increasingly becoming resistant to conventional antibiotics, it would be desirable to provide alternative and preferably prophylactic means of protecting against and fighting microbial infection.

Meningococcal meningitis is a major problem worldwide and in many countries the incidence of infection is increasing. *Neisseria meningitidis* is the causative agent of the disease and is also responsible for meningococcal septicaemia, which is associated with rapid onset and high mortality, with around 22% of cases proving fatal.

At present, vaccines directed at providing protective immunity against meningococcal disease provide only limited protection because of the many different strains of *N. meningitidis*. Vaccines based upon the serogroup antigens, the capsular polysaccharides, offer only short lived protection against infection and do not protect against many strains commonly found in North America and Europe. A further drawback of these vaccines is that they provide low levels of protection for children under the age of 2 years, one of the most vulnerable groups that are commonly susceptible to infection.

Gold *et al.* (Journal of Infectious Diseases, volume 137, no. 2, February

1978, pages 112-121) have reported that carriage of *N. lactamica* may assist in the development of natural immunity to *N. meningitidis* by induction of cross-reactive antibodies. This conclusion was based on the observation of cross-reacting antibodies having complement-dependent bactericidal activity produced in response to *N. lactamica* infection. However, Cann and Rogers (J. Med. Microbiol., volume 30, 1989, pages 23-30) detected antibodies to common antigens of pathogenic and commensal neisseria species, but observed also that antibody to the same antigens was present in both bactericidal and non-bactericidal sera. Thus, it did not seem possible to identify the basis for any cross-reactive bactericidal antibodies, if such exist.

Live attenuated vaccines for meningococcal disease have been suggested by Tang *et al.* (Vaccine 17, 1999, pages 114-117) in which a live, attenuated strain of *N. meningitidis* could be delivered mucosally. Tang also commented on the use of commensal bacteria to protect against infection by pathogenic bacteria, concluding that the cross-reactive epitopes that induce protection against meningococcal infection have not been defined, and therefore that use of genetically modified strains of *N. meningitidis* would be preferred.

It is desirable to provide a further vaccine that gives protective immunity to infection from *N. meningitidis*. It further is desirable to provide a vaccine that confers protective immunity to infants as well as adults and whose protection is long term. It would be of advantage to provide a vaccine that protects against sub-clinical infection, i.e. where symptoms of meningococcal infection are not immediately apparent and the infected individual may act as a carrier of the pathogen. It would further be of advantage to protect against all or a wide range of strains of *N. meningitidis*. It is still further desirable to provide a vaccine against other Neisserial infection, notably gonorrhoea.

It is an object of the present invention to provide compositions containing

immunostimulating components, and vaccines based thereon, that meet or at least ameliorate the disadvantages in the art.

5 The present invention is based on the use of, a commensal *Neisseria* as a vaccine against disease. Accordingly, a commensal species of *Neisseria* such as *N. lactamica* may be used as a live vaccine or a killed whole cell vaccine, or in a vaccine containing fractions of *N. lactamica*. It has surprisingly been demonstrated that mice immunised according to the present invention with *N. lactamica* killed whole cells and outer membrane preparations are
10 protected from lethal intraperitoneal meningococcal challenge, and that vaccines composed of a detergent extract of *N. lactamica* cells or fractions of this, separated by preparative electrophoresis, also protect mice from lethal meningococcal challenge. These results have been obtained using mice and the mouse model used is regarded as predictive of corresponding
15 immunogenic and vaccinating effects in humans.

Accordingly, a first aspect of the present invention provides a composition for vaccination, comprising a commensal *Neisseria* or an immunogenic component, extract or derivative thereof and a pharmaceutically acceptable
20 carrier.

The composition of the invention is particularly suited to vaccination against infection of an animal. The term "infection" as used herein is intended to include the proliferation of a pathogenic organism within and/or on the
25 tissues of a host organism. Such pathogenic organisms typically include bacteria, viruses, fungi and protozoans, although growth of any microbe within and/or on the tissues of an organism are considered to fall within the term "infection".

30 Commensal organisms are those that coexist in an environment with another organism, such coexistence being beneficial to at least one of the organisms and generally not detrimental to either. A number of different commensal

Neisseria are suitable for use in the invention, and these commensal *Neisseria* may be selected from the group consisting of *N. lactamica*, *N. cinerea*, *N. elongata*, *N. flavescens*, *N. polysaccharea*, *N. sicca* and *N. subflava*. Different species of these commensal organisms are known to colonise the buccal or nasal areas and hence each species may be administered according to the known area of the body it normally colonises. Hence also, use of a composition of the invention may result in stimulation of production of protective antibodies *de novo* or if the patient has already been colonised to a certain extent may result in an enhancement of naturally-existing antibodies.

The term "derivative" is used to describe types and strains of commensal *Neisseria* that are modified or attenuated in some way so as to differ from the wild type species. For example, a vaccine composition comprising a recombinant commensal *Neisseria* that exhibits resistance to certain types of antibiotic compounds might advantageously be utilised in combination with such antibiotics in the treatment of infection.

It is an advantage of the invention that vaccination against neisserial diseases may thus be achieved using a non-pathogenic species of *Neisseria*, rendering the vaccination a safer procedure. Furthermore, the protection conferred surprisingly may not be restricted to a specific serotype, subtype or serogroup of the meningococcus but is of general protective efficacy.

A further advantage of the invention is that the commensal *Neisseria* that are the subject of the invention can not revert to virulent types, which can be a risk in the use of live, attenuated strains of virulent bacteria. It is known in the vaccination field to use live, attenuated pathogens and this use carries the risk that the attenuated organism may revert to virulence. This risk is avoided by the present invention. Furthermore, *N. meningitidis* possesses many virulence factors the precise roles of which in pathogenesis are unknown and may possess hitherto unrecognised virulence factors.

Therefore, an additional advantage of the invention is that a composition of the invention can be used with confidence in its level of safety.

5 The method of the invention is of application to vaccination against various infections, preferably to neisserial infections. In a specific embodiment of the invention, protection against meningococcal disease, more specifically caused by *N. meningitidis*, has been demonstrated. The invention is also of application to vaccination generally against neisserial infection, including gonorrhoeal infection, and also to infection from other pathogenic microbial
10 organisms. The invention further provides for vaccination that is aimed at either stimulating or desensitizing the immune system.

The composition can specifically comprise killed commensal *Neisseria*, which may for example be obtained by heat or by suspending commensal *Neisseria*
15 in a mixture of bactericidal agents such as thiomersal and formaldehyde.

The composition may also comprise live commensal *Neisseria*. As mentioned it is usually not required to use attenuated commensal *Neisseria* as these organisms are avirulent.
20

In an embodiment of the invention, an immunogenic component or extract of a commensal *Neisseria* is selected from an outer membrane vesicle preparation of, an outer membrane preparation of, lipooligosaccharide and a protein fraction.
25

The outer membrane preparation and protein fraction of *N. lactamica*, for example, can be obtained from *N. lactamica* cultured in the presence or absence of iron. The protein fraction of *N. lactamica* is conveniently obtained by suspending *N. lactamica* cells or membranes in the presence of detergent
30 and incubating the suspension so as to extract proteins from the *N. lactamica*.

Alternatively, a number of other techniques are known for extraction of outer membrane components - such as protein fractions, lipooligosaccharides and lipopolysaccharides - from cell preparations and are suitable to obtain the commensal *Neisseria* immunogenic components or extracts of the invention.

5 Examples of conventional techniques for this purpose include the use of variation in salt concentration, chaotropic agents, variation in pH (high or low), enzymic digestion and mechanical disruption.

10 A number of different fractions are suitable for use in vaccinating against meningococcal disease. Particularly suitable fractions are those of molecular weight less than 40kDa, of molecular weight more than 40kDa and less than 67kDa, and of molecular weight more than 67kDa.

15 In more specific embodiments of the invention there is provided a composition for eliciting an immune response and suitable for use in vaccinating an individual against Neisserial infection, more specifically meningococcal disease, comprising an antigenic component or antigenic components having the properties:-

- 20 (a) molecular weight 50kDa or lower;
(b) obtainable from *N. lactamica*; and
(c) antibodies to the component(s) obtained from *N. lactamica* cross-react with *N. meningitidis*.

25 In use of a composition containing such a component, extracted using detergent, all mice treated with this component survived a challenged dose of 2×10^7 CFU *N. meningitidis* and three out of five mice survived a higher challenge dose of 6×10^8 CFU.

30 Another specific embodiment of the invention lies in a composition for eliciting an immune response and suitable for use in vaccinating an individual against Neisserial infection, more specifically meningococcal disease, comprising an antigenic component or antigenic components having the

properties:-

- (a) molecular weight at least 40kDa and up to 90kDa;
- (b) obtainable from *N. lactamica*; and
- (c) antibodies to the component(s) obtained from *N. lactamica* cross-react with *N. meningitidis*.

In use of such a component of the invention, obtained using a detergent extract of *N. lactamica*, four out of five mice treated with the component survived a challenge dose of 2×10^7 CFU *N. meningitidis* and mice receiving a higher challenge dose of 6×10^8 CFU survived longer than a control group.

A still further embodiment of the invention lies in a composition for eliciting an immune response and suitable for use in vaccinating an individual against Neisserial infection, more specifically meningococcal disease, comprising an antigenic component or antigenic components having the properties:-

- (a) molecular weight at least 70kDa;
- (b) obtainable from *N. lactamica*; and
- (c) antibodies to the component(s) obtained from *N. lactamica* cross-react with *N. meningitidis*.

In use of such a component, obtained using a detergent extract, one out of five mice survived a challenge dose of 2×10^7 CFU *N. meningitidis* and, whilst all mice succumbed to a higher challenged dose of 6×10^8 CFU, their survival time was longer than a control group which did not receive the component.

In an example of the invention in use, described in more detail below, proteins in the size ranges of 25-35 kDa and 35-43 kDa, extracted from a commensal *Neisseria*, conferred a significant level of immune protection when administered to mice as a vaccine composition.

By way of example of a method of extracting an antigenic component of the

invention, an extraction method comprises:-

- (i) suspending *N.lactamica* cells in an aqueous solution of detergent;
- (ii) incubating the suspension so as to extract the antigenic component from the *N.lactamica*;
- (iii) centrifuging the suspension to separate the suspension into a supernatant and a pellet; and
- (iv) fractionating the antigenic component from the supernatant.

This specific method can be modified according to the extraction protocol selected by the user, for example by using high salt concentration in the initial step (i). In further embodiments of the invention the antigenic component is obtained using recombinant technology by expression of a *N. lactamica* sequence in a suitable host such as *E. coli*.

In a second aspect of the invention there is provided a composition for vaccination against Neisserial infection comprising a commensal *Neisseria* and a pharmaceutically acceptable carrier, wherein the commensal *Neisseria* comprises and expresses a gene from a pathogenic *Neisseria*.

This aspect of the invention offers the benefit of use of a commensal organism to deliver and/or present to the recipient an antigen from a pathogenic *Neisseria*. The gene optionally encodes a surface antigen or a protein that is secreted, and may code for an antigen from, for example, *N. meningitidis* or *N. gonorrhoeae*. The commensal *Neisseria* can be live or killed.

In an embodiment of the second aspect of the invention there is provided a

composition for vaccination against meningococcal disease comprising a commensal *Neisseria* and a pharmaceutically acceptable carrier, wherein the commensal *Neisseria* comprises and expresses a *N. meningitidis* gene.

5 The *N. meningitidis* gene may encode a transferrin binding protein, a SOD for example a Cu,Zn-superoxide dismutase ("SOD"), Neisserial surface protein A ("NspA"), a porin or another outer membrane protein. Gene sequences for the majority of these antigens are known in the literature. Kroll *et al.* in Microbiology 141 (Pt 9), 2271-2279 (1995) describe the
10 sequence of Cu,Zn-SOD. Martin *et al.* in J Exp Med, 1997, April 7th, 185(7), pp1173-1183 describe the sequence of NspA from *N. meningitidis*.

The invention also provides a pharmaceutical composition comprising a composition according to the first or second aspect of the invention plus a
15 pharmaceutically acceptable carrier.

In a third aspect, the invention provides a method of vaccination against Neisserial infection, comprising administering an effective amount of a composition according to the first and second aspects of the invention.

20 In use of an embodiment of the invention described in an example below, there is provided a method of vaccination against meningococcal disease, comprising administering an effective amount of a composition according to the first and second aspects of the invention.

25 In a fourth aspect of the invention there is provided a strain of a commensal *Neisseria*, such as *N. lactamica*, genetically modified so as to express a gene from a pathogenic *Neisseria*. The *N. meningitidis* gene may code for a protein selected from a transferrin binding protein, a SOD for example a Cu,Zn-SOD,
30 NspA, a porin or another outer membrane protein.

The invention further provides, in a fifth aspect a method of extracting a

protein for incorporation in a composition suitable for vaccinating against meningococcal disease, comprising:-

- (i) suspending *N.lactamica* cells in the presence of detergent; and
- (ii) incubating the suspension so as to extract a protein fraction from the cells.

The protein fraction can suitably be of molecular weight 50kDa or lower, at least 40kDa and up to 90kDa or at least 80kDa.

The composition may be combined with a pharmaceutically acceptable carrier - for example the adjuvant alum although any carrier suitable for oral, intravenous, subcutaneous, intraperitoneal intramuscular, intradermal or any other route of administration is suitable - to produce a pharmaceutical composition for treatment of meningococcal disease. Commensal *Neisseria* that are buccal colonizers can be administered in a mouthwash and nasal colonizers in a nasal spray.

Transferrin binding proteins are known to be located on the outer membranes of a number of Gram negative bacteria such as *N. meningitidis*. Formulations of the composition of the present invention with conventional carriers or adjuvants and optionally further supplemented by one or more antigens from *Neisseria* species, optionally recombinantly produced, for example, Cu-Zn SOD, the 22kD NspA, porins, gonorrhoeal antigens or transferrin binding proteins provide a composition for treatment of infection by these bacteria.

In the present invention, the term "transferrin binding protein" or "Tbp" refers to a protein which either alone binds to transferrin or can be part of a complex of proteins that binds transferrin. The term also embraces fragments, variants and derivatives of such a protein provided that

antibodies raised against the fragment, variant or derivative bind the protein. Thus, TbpA and TbpB either dissociated or associated into a complex are considered to be Tbp. Moreover, mutants, fusion proteins or fragments of either TbpA or B or other derivatives of the TbpA + B complex with a common antigenic identity are also considered to be represented by the term Tbp in the present invention.

A live vaccine according to the present invention may be administered via intranasal inoculation. A killed bacteria or subunit vaccine may also be given by this route, or formulated for oral delivery. It is most likely that a subunit vaccine would be administered via the parenteral route. Different commensal *Neisseria* and different strains of *N. lactamica* from those tested in specific embodiments of the invention exist, and the invention is of application also to those other strains.

A sixth aspect of the invention provides a composition comprising an antibody, wherein the antibody binds to a commensal *Neisseria* or an immunogenic component or extract thereof. In use, the antibody can be formulated into a pharmaceutical composition for treatment of Neisserial infection, such as meningococcal disease or infection caused by other *Neisseria*.

An antibody according to this aspect of the invention can be obtained following standard techniques by inoculating an animal with a commensal *Neisseria* or an immunogenic component or extract thereof and thereafter isolating antibodies that bind to the commensal *Neisseria* or the immunogenic component or extract thereof.

A further aspect of the invention provides for a composition comprising a commensal *Neisseria*, or an immunogenic component, extract or derivative thereof, wherein said *Neisseria* comprises a heterologous gene product.

Heterologous gene products of the invention typically include peptides, proteins and antisense sequences that are coded for by a gene sequence that is not native to the commensal *Neisseria*. Typical heterologous gene products of the invention include, for example, bacterial proteins, viral surface peptides, antigens and antibodies and fragments thereof. The heterologous gene product of the invention may be typically any antigen found in a pathogenic organism.

In an embodiment of the invention, the composition comprises a commensal *Neisseria* into which has been transformed an expression vector containing a gene sequence encoding a heterologous gene product. Specific proteins suitable for use in the invention typically include:-

Viral proteins - such as hepatitis B virus surface antigen; rabies virus glycoprotein G; herpes simplex virus glycoprotein D; Epstein-Barr virus glycoprotein; influenza virus glycoprotein; vesicular stomatitis virus nucleoprotein; human respiratory syncytial virus glycoprotein G; human immunodeficiency virus (HIV) envelope; rotavirus subunits; measles virus subunits; and vaccinia virus subunits.

Bacterial proteins - such as *Bordetella pertussis* fimbrial subunits; *Bordetella pertussis* surface proteins; *Bacillus anthracis* subunits; *Escherichia coli* subunits; and *Yersinia pestis* subunits.

Protozoan proteins - such as *Plasmodium falciparum* proteins; trypanosome proteins; and *Cryptosporidium* proteins.

In a further embodiment the composition of the invention suitably provides for a commensal *Neisseria* that expresses a heterologous gene product which is immunostimulatory for treatment of non-infectious disease, for example allergy and cancer. In an example of the invention in use a commensal *Neisseria* that expresses peanut antigens is used to desensitize

a patient with acute peanut allergy.

5 The invention further provides for a commensal *Neisseria* that is transformed with an expression vector that comprises a signal sequence that directs the heterologous gene product to the outer membrane of the *Neisseria* cell. Other signal sequences are also suitable for use in the invention, such as secretion signals or cellular subcompartment localisation signals e.g. periplasmic localisation signals.

10 Further aspects of the invention provide methods for preparing compositions. Such methods are suitable for preparing vaccine compositions that elicit protective immunity to microbial infection when administered to an animal.

15 An example of the invention in use, described in more detail below, provides for a method of preparing a composition comprising the steps of:

- a) inserting a gene coding for a heterologous gene product into an expression vector;
- 20 b) transforming said expression vector into a commensal *Neisseria* so that said heterologous gene product is expressed in said *Neisseria*; and
- 25 c) combining the *Neisseria* of (b) with a pharmaceutically acceptable carrier.

A further example of the invention, provides for a method of preparing a composition comprising the steps of:

- 30 a) inserting a gene coding for a heterologous gene product into an expression vector;

b) transforming said expression vector into a commensal *Neisseria* so that said heterologous gene product is expressed in said *Neisseria*;

5 c) obtaining an immunogenic component or extract from the *Neisseria* of (b); and

d) combining the immunogenic component or extract of (c) with a pharmaceutically acceptable carrier.

10

In yet a further example of the invention in use is provided a method of preparing a composition comprising the steps of:

15 a) obtaining an immunogenic component or extract from a commensal *Neisseria*; and

b) combining the immunogenic component or extract of (a) with a heterologous gene product and a pharmaceutically acceptable carrier.

20

Thus, the invention provides for (a) methods and compositions in which an extract is taken from a commensal *Neisseria* that expresses a heterologous gene product, and (b) methods and compositions where an extract is obtained from a commensal *Neisseria* and the heterologous gene product expressed elsewhere (in another organism) is then added to this latter extract.

25

Further aspects of the invention provide for use of a commensal *Neisseria* in the manufacture of a medicament for treatment of *Neisseria* infection, and for use of a commensal *Neisseria*, or an immunogenic component, extract or derivative thereof, wherein said *Neisseria* comprises a heterologous gene product, in the manufacture of a medicament for the treatment of infection

30

or for immunostimulation in an animal.

Specific embodiments of the invention are discussed in more detail by means of the Examples described below. The results referred to in the Examples are illustrated by the accompanying drawings, in which:

Fig.1 shows protection of mice against intraperitoneal ("IP") infection with *N.meningitidis* strain K454 by use of *N.lactamica* whole cells and outer membrane fractions;

Fig.2A shows protection of mice against IP infection with *N.meningitidis* strain K454 by use of detergent and high, medium and low molecular weight extracts of *N.lactamica* cells - upper panel = challenge by 2×10^7 CFU, lower panel = challenge by 6×10^8 CFU;

Fig.2B shows the components of the high, medium and low molecular weight fractions of fig.2A;

Fig.3 shows an immunoblot illustrating cross-reaction of antibodies in sera from meningococcal disease patients with proteins from *N.lactamica* strain Y92-1009;

Fig. 4 shows a photograph of a gel on which subfractions of low molecular weight outer membrane protein extract have been run; and

Fig. 5 shows protection of mice against IP infection with *N.meningitidis* strain K454 when immunised with low molecular weight subfractions - upper panel = challenge by 5×10^6 CFU, lower panel = challenge by 1×10^8 CFU.

Example 1

Preparation of vaccine containing killed whole cells

5 *Neisseria lactamica* strain Y92-1009 was grown in Mueller Hinton broth (MHB) containing $5\mu\text{gml}^{-1}$ ethylenediamine-di(o-hydroxyphenylacetic acid) (EDDHA), incubated at 37°C with shaking (140rpm) for approximately 6h.

10 Bacteria were then harvested by centrifugation and resuspended in phosphate buffered saline (PBS) containing 1% (v/v) formaldehyde and 0.1% (w/v) thiomersal, and left to stand overnight at $2-8^{\circ}\text{C}$. Killed cells were then resuspended in PBS to an OD_{650} of 1.0 (equivalent to $2 \times 10^9 \text{ CFUml}^{-1}$) and alhydrogel added to 25% (V/V), yielding a product suitable for subcutaneous administration.

15 This method is suitable also for *N. cinerea*, *N. elongata*, *N. flavescens*, *N. polysaccharea*, *N. sicca* and *N. subflava*.

Example 2

20 **Preparation of vaccine containing *N. lactamica* outer membrane (OM) preparations**

25 *N. lactamica* strain Y92-1009 was grown in MHB with and without the addition of $5\mu\text{gml}^{-1}$ EDDHA overnight at 37°C with shaking. Iron limited (with EDDHA) and iron replete cells were then treated separately. Bacteria from 1.5 litres were harvested by centrifugation and resuspended in 20ml 200mM Lithium acetate, 5mM EDTA, pH 6.0 and incubated for 3h at 37°C with shaking. Bacteria were then passed 7 times through a 21 gauge needle and pelleted at 8000g for 10min.

30 The supernatant was recovered and membranes pelleted by centrifugation at 100,000g for 1h at 4°C . The membranes were then resuspended in 10mM HEPES, pH 7.4, containing 0.1% (v/v) 10mM PMSF, yielding OM-

containing vaccinating preparations. The protein content of the OM vaccine preparations was determined using the bicinchoninic acid assay (Sigma, UK). OMs were diluted in sterile deionized water to give a protein concentration of $100\mu\text{gml}^{-1}$. This was then mixed with an equal volume of Freund's
5 adjuvant, to give a final protein concentration of $50\mu\text{gml}^{-1}$, and emulsified thoroughly. Freund's complete adjuvant was used for the primary dose, and Freund's incomplete for subsequent boosts.

Example 3

Preparation of vaccine containing lipooligosaccharide (LOS)

Purification of LOS was carried out from *N. lactamica* strain Y92-1009 using the method of Gu, X-X and Tsai, C.M. (1991) Anal Biochem. **196**; 311-318.
15 Vaccine was prepared using Freund's adjuvant as above with LOS at a final concentration of $10\mu\text{gml}^{-1}$.

Example 4

Vaccination and challenge schedule

Groups of 5 mice were vaccinated with each preparation as follows:-

Prime:- Day 0

First boost:- Day 21

25 Second boost:- Day 28

Mice vaccinated with killed cells of Example 1 received 0.5ml subcutaneously, equivalent to 1×10^9 CFU. Mice vaccinated with OM of Example 2 and LOS of Example 3 received 0.2ml subcutaneously; equivalent
30 to $10\mu\text{g}$ of protein and $2\mu\text{g}$ of LOS.

On day 35, mice were challenged by intraperitoneal injection with

approximately 10^8 CFU *N. meningitidis* K454 made up in MHB containing transferrin at a final concentration of 20mg/ml. The mice were then examined and the number of survivors noted and the results are shown in fig.1. After 4 days all 5 mice survived in the groups vaccinated with whole cells and OMPs (with iron) and 3 survived in the group vaccinated with OMPs (without iron). After 5 days all members of the control group and of the group vaccinated with LOS (marked LPS on the figure) had died.

Example 5

Preparation of vaccine comprising *N. lactamica* fractions

Brain heart infusion agar plates were inoculated with 50 μ l of *N. lactamica* strain Y92-1009 and incubated overnight at 37°C, with 5% CO₂. This was used to inoculate a 100ml MHB starter culture which was incubated with shaking at 37°C for 6 h. Starter culture (15ml) was added to each of 6x500ml volumes of MHB. These were then incubated with shaking overnight at 37°C and the conditions were made iron-limited by the addition of 5 μ gml⁻¹ EDDHA. The cells were harvested by centrifugation and the supernatant discarded. The cells were washed with 100ml PBS and then pelleted by centrifugation. Cell pellets were resuspended in PBS + 0.3% (v/v) Elugent (Calbiochem, 2ml per g wet weight) and incubated with shaking at 37°C for 20 min. The cells were then removed by centrifugation and the pellet discarded. EDTA and N-lauroyl sarcosine were then added to the supernatant to 10mM and 0.5% (w/v) respectively.

The BIORAD Prep Cell, model 491 was then used to separate the proteins contained in the detergent extract. A 4cm, 7% acrylamide native resolving gel was cast with a 2 cm stacking gel. 12mg of protein in native sample buffer was electrophoresed using running buffer containing 0.1% (w/v) SDS, 0.025M Tris and 0.192M glycine at 40mA and 400V until the dye front reached the bottom of the gel. 3ml fractions of the eluted proteins were then

collected. Once the fractions were collected they were pooled into groups consisting of proteins of molecular weight approximately less than 40kDa, between 40 and 67kDa and more than 67kDa. The pooled proteins were concentrated by ammonium sulphate precipitation and dialysed against PBS. These were diluted in PBS to a protein concentration of 100ug/ml and Freund's complete adjuvant was added at a ratio of 1:1(v/v) or Freund's incomplete adjuvant for booster doses.

Example 6

Vaccination and challenge schedule

Groups of 5 mice were vaccinated with each preparation as follows:-

Prime:-	Day 0
First boost:-	Day 21
Second boost:-	Day 28

Mice were vaccinated with no vaccine (i.e control group), Elugent ("Registered Trade Mark") extract or high, medium or low molecular weight fraction. The mice receiving the protein fraction groups received 0.2ml subcutaneously; equivalent to 10µg of protein.

On day 35 mice were challenged by intraperitoneal injection with either approximately 2×10^7 or 6×10^8 CFU *N. meningitidis* K454 made up in MHB containing transferrin at a final concentration of 20mg/ml. The mice were then examined over four days and the number of survivors noted, and the results are shown in fig.2A - upper panel 2×10^7 challenge and lower panel 6×10^8 challenge. The components of the high, medium and low molecular weight fractions are shown in fig.2B. after being run an SDS-PAGE gel.

Example 7

We investigated samples of human sera following meningococcal disease and these showed that antibodies were produced which react with a range of *N. lactamica* proteins. The results of the immunoblot are shown in fig.3.

5 **Example 8**

Due to the level of protection offered by the low molecular weight pool in Example 6 (see Fig. 2A), further separation of these proteins was carried out, according to the method of Example 5, to further characterise components responsible for protection. Proteins were pooled into three groups consisting of <25 kDa (g1), 25-35 kDa (g2) and 35-43 kDa (g3) (shown in Fig. 4). Determination of the levels of lipopolysaccharide (LPS) revealed high levels of LPS in fraction g1 [26 580 endotoxin units per ml (EUml⁻¹)], and considerably lower levels in the remaining fractions (9149 EUml⁻¹ in g2 and 9348 EUml⁻¹).

As in previous examples, groups of five mice were immunised, using a three dose schedule with one of the three groups of proteins described above, proteins >43 kDa and detergent extract of killed whole *N. lactamica* cells and killed whole *N. lactamica*. Animals were challenged with *N. meningitidis* serogroup B, strain K454, at a dose of 5×10^6 or 1×10^8 CFU, together with unimmunised controls. The number of survivors on each day post challenge is shown in Fig. 5.

All mice, apart from the control group and one mouse in group g3, survived the lower challenge dose; however, at the higher challenge dose the g2 and g3 protein groups (25-35 kDa and 35-43 kDa respectively) offered best protection.

30 **Example 9**

Commensal *Neisseria* as a vehicle for recombinant protein expression

The gene encoding the measles virus nucleocapsid protein was cloned into the pMGC18.1 shuttle vector (Webb *et al.*, 1998, poster at the 11th International Pathogenic Neisseria Conference, Nice) and transformed into *E.coli* DH5alpha. Expression of the measles virus nucleocapsid protein was confirmed by western blotting probed with specific antiserum. This construct was then used to transform *N. lactamica* by conjugation. Expression of the measles virus nucleocapsid protein was placed under the control of the neisserial *frpC* promoter and expression at high levels was seen when the bacteria were grown under iron-limited growth conditions.

The invention thus provides immunogenic compositions and vaccines for use in protecting against disease.

CLAIMS

1. A vaccine composition comprising a commensal *Neisseria* or an immunogenic component, extract or derivative thereof.
2. A composition according to Claim 1 comprising killed *Neisseria*.
3. A composition according to Claim 2 wherein killed *N.lactamica* are obtained by suspending *N.lactamica* in a mixture of bactericidal agents such as thiomersal and formaldehyde.
4. A composition according to Claim 1 comprising live *N. lactamica*.
5. A composition according to Claim 1 wherein the immunogenic component or extract of *Neisseria* is selected from an outer membrane preparation, a lipooligosaccharide and a protein fraction.
6. A composition according to Claim 5 wherein the *Neisseria* is *N. lactamica*.
7. A composition according to Claim 5 wherein the immunogenic component or extract comprises a protein fraction of molecular weight less than 40kDa.
8. A composition according to Claim 5 wherein the immunogenic component or extract comprises a protein fraction of molecular weight more than 40kDa and less than 67kDa.
9. A composition according to Claim 5 wherein the immunogenic component or extract comprises a protein fraction of molecular weight more than 67kDa.
10. A composition according to any of Claims 7 to 9 wherein a protein

fraction of *N.lactamica* is obtained by suspending *N.lactamica* cells in the presence of detergent and incubating the suspension so as to extract proteins from the *N.lactamica*.

11. A composition for eliciting an immune response and suitable for use in vaccinating an individual against Neisserial infection, comprising an antigenic component having the properties:-
 - (a) it is of molecular weight 50kDa or lower;
 - (b) it is obtainable from a commensal *Neisseria*; and
 - (c) antibodies to the component obtained from the commensal *Neisseria* cross-react with *N. meningitidis*.
12. A composition for eliciting an immune response and suitable for use in vaccinating an individual against Neisserial infection, comprising an antigenic component having the properties:-
 - (a) it is of molecular weight at least 40kDa and up to 90kDa;
 - (b) it is obtainable from a commensal *Neisseria*; and
 - (c) antibodies to the component obtained from the commensal *Neisseria* cross-react with *N. meningitidis*.
13. A composition for eliciting an immune response and suitable for use in vaccinating an individual against Neisserial infection, comprising an antigenic component having the properties:-
 - (a) it is of molecular weight at least 70kDa;
 - (b) it is obtainable from a commensal *Neisseria*; and
 - (c) antibodies to the component obtained from the commensal *Neisseria* cross-react with *N. meningitidis*.
14. A composition according to any of Claims 11 to 13 further comprising one or more of a transferrin binding protein, a Cu,Zn-SOD, a porin and NspA.
15. A composition for vaccination against Neisserial infection comprising a

commensal *Neisseria* and a pharmaceutically acceptable carrier, wherein the commensal *Neisseria* expresses a gene from a pathogenic *Neisseria*.

16. A composition according to Claim 15 wherein the commensal *Neisseria* expresses a gene which encodes a protein from *N. meningitidis* selected from the group consisting of transferrin binding protein; a Cu,Zn-SOD; an NspA; a porin; and an outer membrane protein.
17. A pharmaceutical composition comprising a composition according to any of Claims 1 to 14 plus a pharmaceutically acceptable carrier.
18. A method of vaccination against microbial infection, comprising administering an effective amount of a composition according to any of Claims 1 to 17.
19. A method of vaccination against Neisserial infection, comprising administering an effective amount of a composition according to any of Claims 1 to 17.
20. A commensal *Neisseria*, which expresses a gene from a pathogenic *Neisseria*.
21. A commensal *Neisseria* according to Claim 20, which expresses a *N.meningitidis* gene which codes for a protein selected from the group consisting of: a transferrin binding protein, a porin, NspA, an outer membrane protein, and a Cu,Zn-SOD.
22. A method of extracting a protein for incorporation in a composition suitable for vaccinating against meningococcal disease, comprising:-
 - (i) suspending commensal *Neisseria* cells in the presence of detergent; and

- (ii) incubating the suspension so as to extract a protein fraction from the cells.
23. A method according to Claim 22, wherein the protein fraction is of molecular weight 50kDa or lower.
 24. A method according to Claim 22, wherein the protein fraction is of molecular weight at least 40kDa and up to 90kDa.
 25. A method according to Claim 22, wherein the protein fraction is of molecular weight at least 80kDa.
 26. A method according to any of Claims 22 to 25 further comprising mixing the protein fraction with one or more of a transferrin binding protein, a Cu,Zn-SOD, a porin, an NspA and a meningococcal protein.
 27. A pharmaceutical composition comprising an antibody that binds to a commensal *Neisseria* or an immunogenic component or extract thereof and a pharmaceutically acceptable carrier.
 28. A composition for vaccinating against meningococcal disease substantially as hereinbefore described.
 29. A composition comprising a commensal *Neisseria*, or an immunogenic component, extract or derivative thereof, wherein said *Neisseria* comprises a heterologous gene product.
 30. A composition according to Claim 29 wherein the heterologous gene product is a product of a gene that is expressed in said commensal *Neisseria*.
 31. A composition according to Claim 29 wherein the heterologous gene

product is physically combined with said commensal *Neisseria*.

32. A method of preparing a composition comprising:

- a) inserting a gene coding for a heterologous gene product into an expression vector;
- b) transforming said expression vector into a commensal *Neisseria* so that said heterologous gene product is expressed in said *Neisseria*; and
- c) combining the *Neisseria* of (b) with a pharmaceutically acceptable carrier.

33. A method of preparing a composition comprising the steps of:

- a) inserting a gene coding for a heterologous gene product into an expression vector;
- b) transforming said expression vector into a commensal *Neisseria* so that said heterologous gene product is expressed in said *Neisseria*;
- c) obtaining an immunogenic component or extract from the *Neisseria* of (b); and
- d) combining the immunogenic component or extract of (c) with a pharmaceutically acceptable carrier.

34. A method of preparing a composition comprising the steps of:

- a) obtaining an immunogenic component or extract of a


commensal *Neisseria*; and

b) combining the immunogenic component or extract of (a) with a heterologous gene product and a pharmaceutically acceptable carrier.

35. Use of a commensal *Neisseria* in the manufacture of a medicament for treatment of *Neisserial* infection.
36. Use of a commensal *Neisseria*, or an immunogenic component, extract or derivative thereof, wherein said *Neisseria* comprises a heterologous gene product, in the manufacture of a medicament for the treatment of infection in an animal.
37. Use of a commensal *Neisseria*, or an immunogenic component, extract or derivative thereof, wherein said *Neisseria* comprises a heterologous gene product, in the manufacture of a medicament for immunostimulation in an animal.



THIS PAGE BLANK (USPTO)

Fig. 1. Protection of mice against IP infection with *N.meningitidis* strain K454 by vaccination with *N. lactamica* whole cells and outer membrane proteins




THIS PAGE BLANK (USPTO)

Fig. 2A. Protection of mice against IP infection with *N. meningitidis* strain K454 by vaccination *N. lactamica* extracts




THIS PAGE BLANK (USPTO)

Fig. 2B. SDS-PAGE of proteins separated by preparative electrophoresis used in protection expt.



THIS PAGE BLANK (USPTO)


Fig 3. Immunoblot showing cross-reaction of antibodies in sera from meningococcal disease patients with proteins from *N. lactamica* strain Y92-1009



(THIS PAGE BLANK (USPTO))


(-)

Fig.4




THIS PAGE BLANK (USPTO)

Fig.5



No vaccine
 Elugent extract
 Killed whole cells
 g1
 g2
 g3
 Med. + high



THIS PAGE BLANK (USPTO)